1TC 824 C2 A2 no. 69: 64 LIBRARY
UNIVERSITY OF CALIFORNIA
DAVIS







State of California
THE RESOURCES AGENCY

partment of Water Resources

BULLETIN No. 69-64

CALIFORNIA HIGH WATER 1963 - 1964

SEPTEMBER 1965

NUV 1 2 1965

HUGO FISHER

Administrator
The Resources Agency

EDMUND G. BROWN
Governor
State of California

WILLIAM E. WARNE

Director

Deportment of Water Resources

LIBRARY
UNIVERSITY OF CALIFORNIA
DAVIS



State of California THE RESOURCES AGENCY

Department of Water Resources

BULLETIN No. 69-64

CALIFORNIA HIGH WATER 1963 - 1964

SEPTEMBER 1965

HUGO FISHER

Administrator
The Resources Agency

EDMUND G. BROWN
Governor
State of California

WILLIAM E. WARNE

Director

Department of Water Resources

Department of Water Resources

No. 64 . 69-64

CALIFORNIA HIGH WATER

ANY THEMSENDERS

manual a south

-

THE OSUM

FOREWORD

This report, Bulletin 69-64, "California High Water 1963-1964", is the second of an annual series on the subject of high water in California.

This report was prepared to provide (in one report) information on precipitation, and peak flows and stages, resulting from the major storms of the water year. These reports contain a summary of the water year events, a description of the general weather characteristics preceding and during the storm period, a discussion of the precipitation characteristics, a discussion of the resulting runoff, and a review of the flood damage. When significant, tabulations of precipitation comparisons, peak flows and stages, and reservoir storage data are included in the appendixes.

Basic data for these reports are supplied by many governmental and private agencies. Among these are the United States Weather Bureau, United States Geological Survey, United States Bureau of Reclamation, Corps of Engineers, Pacific Gas and Electric Company, East Bay Municipal Utility District, and numerous other public and private districts and agencies.

The data appearing in this report are considered to be accurate and reliable. It should be noted, however, that hydrologic data may be revised (usually the changes are minor) at a later date on the basis of subsequent studies and information. Therefore, all data should be considered to be preliminary and subject to revision.



TABLE OF CONTENTS

			-	Page
FOREWORD	•			iii
LETTER OF TRANSMITTAL			•	vii
ORGANIZATION, DEPARTMENT OF WATER RESOURCES	•		•	viii
CALIFORNIA HIGH WATER, 1963-1964	•			1
Summary	٠	•	•	1
Meteorology of the January 1964 Storm	•	•	•	3
Runoff		•		6
North Coastal Area				6
San Francisco Bay Area	•		•	7
Central Coastal Area			•	8
Central Valley Area	•	•	•	8
Flooded Areas and Flood Damage			•	11
Flood Control Projects Completed 1963-1964	•	•	•	14
TABLE				
Table No.				Page
I SACRAMENTO RIVER FLOOD CONTROL PROJECT, WEIR OVERFLOWS	•	•	•	10
APPENDIX				
APPENDIX A - RUNOFF				
TABLE A-1 PEAK FLOWS AND STAGES	•	•	•	A-1
North Coastal Area	•	•	•	A-1 A-4 A-6 A-7

TABLE OF CONTENTS (Continued)

PLATES

MAJOR DRAINAGE AREAS IN CALIFORNIA	Plate 1
NORTH COASTAL AREA - ISOHYETAL AND STATION LOCATION MAP	Plate 2
SAN FRANCISCO BAY AREA - ISOHYETAL AND STATION LOCATION MAP	Plate 3
CENTRAL COASTAL AREA - ISOHYETAL AND STATION LOCATION MAP	Plate 1
CENTRAL VALLEY AREA - ISOHYETAL AND STATION LOCATION MAP	Plate 5
GAGE HEIGHTS OF SACRAMENTO RIVER AND YOLO BYPASS	Plate (
GAGE HEIGHTS OF FEATHER RIVER, YUBA RIVER, BEAR RIVER, AND CACHE CREEK	Plate 1

PARTMENT OF WATER RESOURCES

MENTO



June 22, 1965

Honorable Edmund G. Brown, Governor and Members of the Legislature of the State of California

Gentlemen:

Bulletin No. 69-64, "California High Water, 1963-1964", is the second of an annual series summarizing water year events. The report provides data on precipitation, streamflows, and flood damage resulting from the major storms of the 1963-64 water year, and is intended for the information of both governmental and private agencies, as well as the general public.

Sincerely yours,

Willi E. Laure

Director

State of California The Resources Agency DEPARTMENT OF WATER RESOURCES

EDMUND G. BROWN, Governor
HUGO FISHER, Administrator, The Resources Agency
WILLIAM E. WARNE, Director, Department of Water Resources

ALFRED R. GOLZE'
DIVISION OF OPERATIONS
ROBIN R. REYNOLDS Division Engineer WILLIAM L. HORN Chief, Water Operations Branch
This report was prepared under the immediate supervision of
ROBERT W. MILLER Chief, Flood Operations & Forecasts Unit
by
WILLIAM A. ARVOLA Meteorologist III
and
CHARLES H. HOWARD Associate Engineer, W.R.
assisted by
RICHARD A. McGUIRE Engineering Aid II
RICHARD SPELLMAN Student Trainee

CALIFORNIA HIGH WATER

1963 - 1964

Summary

The 1963-64 water year (October 1 - September 30) was essentially dry, with only one major period of precipitation intense enough to produce significant runoff.

At the beginning of the water year, the October November period was marked by several storms in which most of
the state received above-average rainfall. Some locations in
southeastern California received 3 to 4 times normal amounts
as a result of these storms. Resultant runoff amounts and
streamflows were generally moderate in Northern and Central
California, but Southern California reported scattered instances
of minor flooding and some damage to property and crops.

In contrast, very little precipitation fell anywhere in the state during the month of December; so little, in fact, that despite the above-normal amounts of the previous months, the accumulated total at the end of December was below normal for most areas.

This trend was reversed during the month of January in the North Coast and central part of the state, and particularly in the Sierra Nevada, with precipitation well above normal at these locations. Elsewhere in California the unseasonably dry weather continued.

As the most significant high water events of the year were associated with the storm period of January 17-27, the meteorology, runoff, and damages associated with this storm are described in detail in this bulletin.

From February to May, the remainder of the usual high water season, unusually dry conditions prevailed in most of California, and resulting streamflows were held to belownormal values. The month of February was established as one of the driest on record, with less than one inch of precipitation at most stations in the state. By May 1, the water year was noted as one of the 10 driest that had occurred in the last 50 years.

Meteorology of the January 1964 Storm

December 1963 in California can be described as having been cold, foggy, and with very little rain. High pressure systems over the Great Basin plateau dominated the circulation pattern, and only a few storm fronts penetrated into California.

In January the Aleutian low pressure center began to deepen and spread out on a west-to-east axis. To the south, the central Pacific anticyclone, also much stronger than normal, extended across the Pacific. The strong pressure gradient (south-to-north) between the two systems stimulated a strong westerly flow through the United States. More weather fronts began to move through California, but the resulting precipitation was not unusually heavy in the first half of the month.

A cold front entered California in the North Coast area on the morning of January 17, and by the next morning had moved into the Los Angeles area. The rainfall from this system was of only moderate intensity. A new storm which developed in the central Pacific south of the Aleutians near the 50th parallel began to move toward the Washington-Oregon coast on the morning of January 18. The frontal system in this storm had the structure of an open wave cyclone with a very wide warm sector, and was located in the north of a fairly intense Pacific high pressure cell centered along the 35th parallel. The warm

front of this system reached the Northern California coast late on the 18th, preceded by an area of precipitation. The freezing level ahead of the warm front in the Central Sierra was about 3,500 feet, and although the passage of the warm front across the state was not clearly marked by observations at various stations, some warming occurred in the mountains with the freezing level lifting to 5,000 feet. Snow fell at elevations above this level. The warm front rains at lower elevations were heaviest in the North Coast area and in the Sierra Nevada, with only light amounts at low elevations in the San Francisco Bay area and the Sacramento Valley.

The cold front of this storm was delayed in entering California until the early morning hours of January 20, passing Eureka at 0100 hours, San Francisco at 1900 hours, and Sacramento at 2000 hours. This cold front was accompanied by heavy precipitation. In the hour ending at 1900 hours, San Francisco International Airport had a one-hour rainfall of 0.87 inch; the maximum six-hour amount was 2.06 inches. Sacramento (City Office, USWB) had two hours with 0.41 inch in each hour; the maximum six-hour amount was 1.97 inches. These precipitation rates represent fairly intense frontal precipitation. The cold front precipitation at low elevations had durations of 6 to 12 hours.

The snow depth at Blue Canyon (elevation 5,280 feet) on Interstate Highway 80 increased from 4 inches on the morning of January 17 to 57 inches on the morning of January 22, although

some of the precipitation which fell during the warmest part of the storm on the 19th fell as rain mixed with snow. The largest accumulation at Blue Canyon occurred during the period from the 20th to the 22nd.

Isohyetal maps for this storm, covering the 10-day period January 15-25, 1964, are shown on Plates 2 - 5. The amounts of maximum storm precipitation were 14 inches in the Smith, lower Klamath, Mattole, Eel, upper Sacramento (above Shasta Dam), Feather, Yuba, and American River Basins. Most of the precipitation in this 10-day period fell during the 5 days beginning on January 17.

Runoff

In general, low to moderate flows occurred in Northern and Central California streams during the period January 20-24, although in several locations new maximum flows of record were established. The following is a brief discussion of the runoff in each basin affected, including comparisons with peaks from the 1962-63 water year, as well as with maximum peaks of record.

North Coastal Area

Most of the northerly basins had moderate flows, slightly below those recorded in 1962. The Smith River near Crescent City had a peak of 93,400 cfs, as compared to 113,000 cfs in 1962, and a maximum recorded peak of 165,000 cfs in 1955. The Klamath River near Klamath had a peak flow of 162,000 cfs, as compared to 176,000 cfs in 1962, and a maximum recorded peak of 425,000 cfs in 1955.

Further south, the Mad River near Arcata recorded 45,200 cfs, compared to 28,900 cfs in 1962, and a maximum of record of 77,800 cfs in 1955. The Eel River at Scotia had a peak flow of 178,000 cfs, as compared to 252,000 cfs in 1963, and a maximum recorded flow of 541,000 cfs in 1955. Several smaller streams in this area had new peak flows of record, as follows: Willow Creek at Willow Creek, 5,260 cfs, as compared to a previous maximum of 5,190 cfs in 1962; Little River at

Crannell, 11,200 cfs, compared to 9,300 cfs in 1957; and the North Fork Mad River near Korbel, 8,400 cfs surpassing 7,170 cfs in 1960.

In the Russian River basin, flows were generally somewhat below those of the previous year. Dry Creek near Geyserville, a tributary of the Russian River, recorded 10,200 cfs, compared to a peak of record of 25,800 cfs established in 1963. The Russian River near Guerneville peaked at 33,400 cfs, compared to 71,800 cfs in 1963, and a maximum of record of 90,100 cfs in 1955.

San Francisco Bay Area

Streamflows in the San Francisco Bay area can be classified in the low to moderate range, with a few isolated high flows, in contrast to 1962-63, when many new peak flows of record were established.

In the North Bay area, the Napa River near Napa had a flow of 5,260 cfs, compared to a maximum of record of 16,900 cfs in 1963. Several creeks in this vicinity had peaks which equalled or exceeded previously established maximums. These were Walker Creek near Tomales with a flow of 3,430 cfs, equalling the record flows of 1961 and 1962, and Novato Creek near Novato with 1,330 cfs, exceeding the previous maximum of 1,190 cfs in 1958.

East Bay, South Bay, and Penninsula area streams had generally low flows, as typified by Walnut Creek at Walnut

Creek, with a peak of 2,760 cfs, compared to 11,400 cfs in 1962, and a maximum of record of 12,200 cfs in 1958.

Central Coastal Area

Low runoff occurred in most streams in this area, considerably below that of the previous year, and at several locations in the vicinity of Santa Maria no flow whatsoever was recorded.

The following are typical examples of the magnitude of flows in the Central Coast basins: the San Lorenzo River at Big Trees peaked at 2,660 cfs, in contrast to 13,000 cfs in 1963, and a maximum of record of 30,400 cfs in 1955. The Pajaro River at Chittenden had a flow of 1,460 cfs, compared to 11,400 cfs in 1963, and a maximum recorded peak of 24,000 cfs in 1955. Arroyo Seco near Soledad recorded 4,850 cfs, compared to 22,800 cfs in 1963, and a maximum of 28,300 cfs in 1958. The Nacimiento River near Bryson had 6,070 cfs, in contrast to 22,700 cfs in 1963, and a peak of record of 30,300 cfs established in 1955.

Central Valley Area

Runoff in the major basins and tributary streams of the Central Valley north of the Cosumnes River ranged from low to moderate; in most locations somewhat less than that of 1962-63. In the remainder of the valley, the San Joaquin River and its tributaries south from the Mokelumne River basin

recorded no increases over normal winter base flows. Plots of gage heights on the Sacramento River and Yolo Bypass are depicted on Plate 6, and those of the Feather, Yuba, and Bear Rivers and Cache Creek are shown on Plate 7.

In the north, Shasta Lake had moderate inflows from the Sacramento and McCloud Rivers and Squaw Creek, with very low inflow recorded from the Pit River. During the 10-day period January 18-28, storage in the reservoir increased approximately 245,000 acre-feet. Below Shasta, the peak discharge return to the Sacramento River at Keswick was 8,020 cfs. Downstream from Keswick, typical examples of flows in the upper valley tributaries are these: on the west side, Cottonwood Creek near Cottonwood had a flow of 13,000 cfs, compared to 23,100 cfs in 1963, and a maximum of 52,300 cfs in 1941; while Thomes Creek at Paskenta recorded 3,390 cfs, compared to 19,200 cfs in 1963, and a maximum of record of 23,500 cfs in 1955. On the east side of the valley, Cow Creek near Millville peaked at 17,900 cfs, compared to 24,800 cfs in 1962, and a maximum of record 45,200 cfs in 1951. Deer Creek near Vina had 7,880 cfs, in comparison to 10,200 cfs in 1962, and a maximum of record of 23,800 cfs in 1937.

In the Feather, Yuba, and American River basins, flows were very low, far below those of the previous year, when many new peak flows of record were established. In the Feather River basin, the Middle Fork Feather River near Merrimac had a flow of only 3,510 cfs, compared to a maximum

peak of 65,400 cfs in 1963. The Yuba River at Englebright Dam recorded 7,760 cfs, compared to a maximum of record of 150,000 cfs in 1963. On the American River, the Middle Fork near Auburn peaked at 3,410 cfs, in contrast to the record of 121,000 cfs in 1963.

Within the Sacramento River Flood Control Project, all flows were well below maximum project design quantities. The Sacramento River at Sacramento recorded a peak of 52,800 cfs, compared to a project design flow of 110,000 cfs at this location. Overflow occurred at Colusa and Tisdale Weirs, as shown on the following table:

TABLE I
Sacramento River Flood Control Project
WEIR OVERFLOWS

Weir	Beginnin Overfl Date		Length of Overflow Period	Peak Stage and Discharge			
Moulton				No	Flow		
Colusa	1/21/64	2045	31 hours	63.81	9,810 cfs		
Tisdale	1/22/64	0115	57 hours	47.59	6,390 cfs		
Fremont				No Flow			
Sacramento				No gates opened			

South of Sacramento, runoff amounts generally tapered off and little or no increase over normal winter base flows were recorded in any of the major streams and tributaries of the San Joaquin Valley.

Flooded Areas and Flood Damage

Flood damage in January 1964 was considerably less than in previous high water years, with minor local flooding occurring in only a few locations.

In the North Coastal area, approximately one hundred residents of Orick were forced to evacuate homes and trailers as Redwood Creek overflowed its banks into low lying portions of the town. At the same time, the Smith River near Crescent City also experienced overbank flow, however neither location reported significant damages. South of Eureka, minor damage in the Eel River delta near Fernbridge and Loleta consisted of bank erosion and deposits of debris. Total damage in the entire area was estimated at \$300,000.

The San Francisco Bay and Russian River areas reported scattered instances of high water damage and inconvenience from flooding due to inadequate drainage facilities. Colma Creek in South San Francisco overbanked and inundated a portion of the Bayshore Freeway, interrupting peak hour commuter traffic. Extensive wind damage was also reported in the Bay area.

In the Central Valley, no serious flooding problems occurred, as precipitation from the storm fell as snow at very low elevations, holding river rises to a minimum. This unusually low snowfall, at some locations lower than 1,000 feet elevation, caused many power and telephone outages and road closures lasting several days or more. In the vicinity of

Sacramento, heavy rainfall caused temporary power failures, disruption of telephone service, and many flooded gutters and intersections as the capacity of drainage systems was exceeded. Near Wheatland, along the Camp Far West Canal south of the Bear River, a construction crew dynamited a levee which had impounded drainage water behind it and was causing flooding of nearby homes.

During the water year, several unusual and disastrous events involving water-caused damage also occurred, and although not directly associated with floods they are mentioned here as a matter of general interest.

The first of these events was the failure of the Baldwin Hills Reservoir, a 900 acre-foot capacity municipal reservoir owned and operated by the Los Angeles Department of Water and Power. On the afternoon of December 14, 1963, the main embankment of the reservoir ruptured, releasing a 10-foot deep wall of water which coursed down a one-half mile long canyon lined with homes and ultimately fanned out through a flat residential district. Five persons were killed, about two dozen homes were completely demolished, several hundred were made uninhabitable by heavy damage and silting, and some 2,000 more suffered lesser damages. The flooded area was about one-half square mile, and the total damage amounted to more than \$15,000,000. Subsequent investigation revealed that the cause of failure was gradual foundation subsidence culminating in a sudden earth movement on the day of the

disaster, allowing flow of water through and ultimate collapse of a portion of the embankment.

The second of these occurrences was in the form of a seismic sea wave (popularly misnamed "tidal wave") which struck the town of Crescent City shortly after midnight on March 28, 1964. This wave was a direct result of the disastrous earthquake which had devastated many communities in Alaska on the afternoon of Good Friday, some eight hours earlier. A wall of water 5 - 10 feet in height swept through the harbor of Crescent City and into a portion of the town, causing widespread damage to a 29 block area. Ten persons lost their lives, and total damage to public and private property amounted to approximately \$11,000,000.

Flood Control Projects Completed 1963-1964

In the Central Valley area, several projects which provide important flood control benefits were completed and placed in operation during the 1963-64 water year. These completed projects are covered below.

Black Butte Reservoir, located about 9 miles west of Orland on Stony Creek, provides flood protection to the towns of Orland and Hamilton City as well as protection to about 64,000 acres of farm land west of the Sacramento River. This reservoir, with a gross capacity of 160,000 acre-feet and a maximum flood control storage reservation of 150,000 acre-feet, is owned and operated by the U.S. Army Corps of Engineers.

Camanche Reservoir on the Mokelumne River, is located about 20 miles northeast of Stockton. It provides flood protection to about 69,000 acres of agricultural land; to 3,000 acres of urban and suburban land; and to the towns of Lodi, Woodbridge, and Thornton. The gross storage capacity of the reservoir is 431,500 acre-feet, with a maximum flood control storage reservation of 200,000 acre-feet. It is owned and operated by the East Bay Municipal Utility District.

New Hogan Reservoir is located on the Calaveras
River about 28 miles northeast of Stockton, with the damsite
immediately below that of the smaller Hogan Dam, which is now
inundated. The project provides flood protection for about
46,000 acres of highly developed agricultural land along the

Calaveras River and about 14,000 acres of urban and suburban land in and adjacent to the City of Stockton. This reservoir is owned and operated by the U. S. Army Corps of Engineers, and has a gross storage capacity of 325,000 acre-feet with a maximum flood control storage reservation of 165,000 acre-feet.

Several units of the Department of Water Resources' Lower San Joaquin River Flood Control Project, which extends from the Merced River to Friant Dam, were also completed in 1963-64. Scheduled for completion in 1967, this project comprises many miles of levees and bypass systems designed to prevent flooding problems along the San Joaquin River and its tributaries in this area.



APPENDIX A

RUNOFF

TABLE A-1 PEAK FLOWS AND STAGES

The sample of th

WW/OWNER

SSHIPTS OUR EXOLUTIONS AND STRINGS

APPENDIX A

TABLE A-1

PEAK FLOWS AND STAGES
(Preliminary Data, Subject to Revision)

	Drainage	Period	Source	ource Previous Maximum of of Record					1963 - 1964				
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs			
North Coastal Area													
Middle Fork Smith River at Gasquet	130	1911-18 1958-	USGS	12/2/62	12.28	26,700	1/20	0200	10.52	18,200			
Smith River near Crescent City	613	1931-	USGS	12/22/55	41.20	165,000	1/20	0200	31.22	93,400			
Shasta River near Yreka	796	*1933-	USOS	12/22/55	9.43	6,090	1/20	1000	7.83	3,730			
Scott River near Port Jones	662	1941-	USGS	12/22/55	21.40	38,500	1/20	1600	9.96	5,860			
Klamath River nr. Seiad Valley	6,980	1912 - 25 1951-	USGS	12/22/55	29.2 ^h	122,000 ^c	1/20	1630	12.05	20,100 ^c			
South Fork Salmon River near Porks of Salmon	252	1957-	USGS	12/22/55	18.86 ^h	24,200	1/20	0530	11.10	8,110			
North Pork Salmon River near Porks of Salmon	205	1958-	USGS	12/2/62	14.27	10,500	1/20	0615	9.69	3,650			
Salmon River at Somesbar	746	*1911-	USGS	12/22/55	28.80	84,000	1/20	0700	11.23	19,300			
Klamath River at Somesbar	8,480	1927-	USGS	12/22/55	59.4 ^h	202,000 ^c	1/20	0900	26.30	57,900°			
Red Cap Creek near Orleans	56.1	1958-	USGS	12/2/62	10.69	5,340	1/20	0500	9.50	3,700			
Bluff Creek near Weitchpec	74.6	1958-	USGS	12/22/55	13.7 ^h	20,200	1/20	0300	10.16	7,500			
Trinity River above Coffee Creek near Trinity Center	149	1957~	USGS	2/24/58 12/22/55	10.50 10.5 h	12,800 12,800	1/20	0600	4.34	1,200			
Trinity River at Lewiston	726	1911-	USGS	12/22/55	27.3	71,600	1/20	1200	4.99	570 ^c			
North Fork Trinity River at Helena	151	1911-13 1957-	USGS DWR	1/12/59	19.66	13,500	1/20	0640	13.49	4,820			
Trinity River nr. Burnt Ranch	1,438	1931-40 1956-	USGS	12/22/55	43.2 ^h	172,000	1/20	1700	13.46	15,800°			
New River at Denny	173	1927-28 1959-	USGS	12/2/62	11.71	9,580	1/20	0700	10.17	6,980			
South Fork Trinity R. at Porest Glen	208	1959-	USGS	12/22/55	25.26 ^h	42,400	1/20	1130	15.68	11,400			

TABLE A-1 (Continued)

Stream and Station	Drainage Area in	Period of	Source	Previous Maximum of Record				1963 - 1964		
	Sq. Mi.	Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
North Coastal Area (Co	ntinued)									
South Fork Trinity River near Hyampom	342	1956-	USGS	12/22/55	22.2 ^{h,b}	39,400	1/20	1200	14.50	13,300
Hayfork Creek near Hayfork	87.2	1956-	USGS	2/8/60	11.67	4,210	1/20	0915	10.21	3,190
Hayfork Creek near Hyampom	379	1953-	USGS	12/22/55	18.00	25,300	1/20	1200	13.98	14,200
South Fork Trinity River near Salyer	899	1911-13 1950-	USGS	12/22/55	39.4 ^h	65,100	1/20	1500	26.00	27,800
Willow Creek at Willow Creek	43.3	1959-	USGS	12/2/62	10.02	5,190	1/20	0600	9.86	5,260**
Trinity River near Hoopa	2,848	*1911-	USGS	12/22/55	36.90	190,000	1/20	1800	21.68	62,300 ^c
Klamath River near Klamath	12,100	*1910-	USGS	12/22/55	49.7 ^h	425,000 ^c	1/20	2300	27.10	162,000 ^c
Redwood Creek at Orick	278	1911-13 1953-	USGS	1/18/53 12/22/55	23.95 ^h 23.95	50,000 50,000	1/20	0800	21.97	37,700
Little River at Crannell	44.3	1955~	usgs	3/11/57	9.96	9,300	1/20	0415	10.83	11,200**
Mad River near Forest Glenn	143	1953-	USGS	12/22/55	24.5 ^h	39,200	1/20	2000	9.93	7,300
North Fork Mad R. near Korbel	40.5	1957-	USGS	2/8/60	16.17	7,170	1/20	0400	15.75	8,400**
Mad River near Arcata	484	1910-13 1950-	USGS	12/22/55	27.30 ^b	77,800	1/20	1000	16.04	45,200°
Jacoby Creek nr. Freahwater	6.07	1954~	USGS	12/30/54	7.20	1,670	1/20	0245	5.03	900
Elk River near Falk	44.2	1957-	USGS	2/14/59	27.62	3,220	1/20	0715	27.13	3,050
Eel River below Scott Dam near Potter Valley	290	1922-	USGS	12/11/37	22.9 ^h	41,100 ^c	1/21	1800	8.20	2,600 ^c
Eel River at Van Aradale Dam near Potter Valley	349	1909-	USGS	12/22/55	31.4 ^h	48,600	1/20	1600	11.70	4,710
Outlet Creek nr. Longvale	162	1956-	USGS	2/8/60	20.27	26,500	1/20	1200	14.31	13,600
Eel River above Doa Rios	705	1950-	us c s	12/22/55	45.4h	123,000 ^c	1/20	1700	23.25	27,800°
Black Butte River near Covelo	162	*1951-	usgs	12/21/55	35.8 ^{h,b}	25,000	1/20	1400	12.93	6,820

TABLE A-1 (Continued)

	Drainage	Period	Source	Previous Maximum of Record			1963 - 1964				
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Diachg. in cfs	Date	Time	Stage in ft.	Diachg. in cfs	
North Coastal Area (Co	ontinued)										
Not all address 112 ca	,										
M. F. Eel River below Black Butte River, nr. Covelo	367	1951-	USOS	12/21/55	25.0 ^h	89,100	1/20	1300	13.53	20,500	
Eel River below Dos Rios	1,484	1911-13 1951-	USGS	12/22/55	49.86	283,000°	1/20	1800	26.13	76,900	
North Fork Eel River near Mina	250	1953-	USGS	12/22/55	24.00	58,400	1/20	1200	19.50	27,800	
Eel River at Alderpoint	2,079	1955-	USGS	12/22/55	72.5 ^h	376,000°	1/20	2100	36.59	100,000 ^c	
South Fork Eel R. nr. Branscomb	43.9	1946-	USGS	12/22/55	16.20	20,100	1/21	1300	10.00	6,860	
Tenmile Creek nr. Laytonville	50.3	1957-	USGS	12/22/55	22.9 ^h	16,300	1/20	1100	13.86	6,760	
South Fork Eel R. near Miranda	537	1939-	USGS	12/22/55	42.7 ^h	173,000	1/20	1700	24.00	59,000	
Bull Creek nr. Weott	28.1	1960-	USGS	2/10/61 1/31/63	16.88 16.12	3,400 4,120	1/20	0300	14.33	1,930	
Larabee Creek near Holmes	84.1	1959-	USGS	2/8/60	12.40	10,000	1/20	0600	11.33	7,890	
Eel River at Scotia	3,113	*1910-	USGS	12/22/55	61.90	541,000	1/21	0200	39.40	178,000	
South Fork Van Duzen River nr. Bridgeville	36.2	*1951-	USGS	12/22/55	11.91 ^{b,h}	8,990	1/20	0600	13,63	6,760	
Van Duzen River nr. Bridgeville	216	1950-	USGS	12/22/55	21.3 ^h	43,500	1/20	0800	18.10	32,000	
Mattole River nr. Petrolia	240	*1911-	USGS	12/22/55	29.60	90,400	1/20	0800	21.85	43,200	
Noyo River nr. Fort Bragg	106	1951-	USGS	12/22/55	25.64	22,000	1/20	unknown	17.72	6,570	
Rancheria Creek near Boonville	65.6	1959-	USGS	1/31/63	18.30	13,900	1/20	1500	12.65	6,920	
Navarro River near Navarro	303	1950-	USGS	12/22/55	40.60	64,500	1/20	2100	25.80	17,900	
South Fork Gualala River nr. Annapolis	161	1950-	USGS	12/22/55	24.57	55,000	1/20	1800	13.60	15,000	
Rusaian River near Ukiah	99.6	*1911-	USGS	12/21/55	21.0	18,900	1/20	1500	13.28	8,850	
East Fork Rusaian River nr. Calpella	93.0	1941-	USGS	12/21/55	15.06 ^b	13,300 ^e	1/20	1600	12.85	8,340°	

TABLE A-1 (Continued)

Stream and Station	Drainage	Period of	Source	Pre	1963 - 1964					
	Area in Sq. M1.	Record	Record (a)	Date	Stage in ft.	Diachg. in cfs	Date	Time	Stage in ft.	Dischg. in cfa
North Coastal Area (Co	ntinued)									
Russian River near Hopland	362	1939-	USGS	12/22/55	27.00	45,000 ^c	1/20	2000	17.72	17,000°
Peliz Creek mear Hopland	31.1	1958-	USGS	1/31/63	13.43.	2,910	1/20	1500	13.60	2,710
•	<i></i>	-22-		12/23/55	13.43 _h 13.60 ^h	2,710	-/	_,,,,	23.00	-,,20
Aussian River ar. Cloverdale	502	1951-	USGS	12/22/55	30.9 ^h	53,000 ^c	1/20	2315	18.23	16,600°
Big Sulphur Cr. mear Cloverdale	82.3	1957-	USGS	12/22/55	22.2 ^h	20,000	1/20	1545	11.21	5,830
Russian River nr. Healdsburg	793	1939-	USGS	2/28/40	30.0	67,000	1/21	0500	14.11	22,100
Ory Creek near Lloverdale	87.8	1941-	USGS	1/31/63	17.91	17,700	1/20	1600	9.55	4,720
Ory Creek nr. Geyserville	162	1959-	USGS	1/31/63	16.50	25,800	1/20	1800	9.78	10,200
Santa Rosa Creek Lear Santa Rosa	12.5	1959-	USGS	2/8/60	13.35 ^h	3,200	1/20	1700	8.95	1,040
Russian River nr. Guerneville	1,340	*1939-	USGS	12/23/55	49.7 ^h	90,100 ^c	1/21	0900	29.88	33,400
ustin Creek mear Cazadero	63.1	1959~	USGS	2/13/62	20.6 ¹	15,100	1/20	1600	12.45	6,150
San Francisco Bay Area	_									
dalker Creek nr. Tomales	37.1	1959-	USGS	1/31/61 2/13/62	18.18 17.72	3,430 3,430	1/20	2000	18,52	3,430
Corte Madera Creek at Ross	18,1	1951-	USGS	12/22/55	17.45	3,620	1/20	1800	10.07	1,010
Novato Creek near Novato	17.5	1946-	USGS	2/24/58	8.24	1,190	1/20	1800	8.74	1,330
Sonoma Creek it Boyes Hot Springs	62.2	1955-	USGS	12/22/55	17.10	8,880	1/20	1800	11.69	4,360
Wapa River nr. St. Helena	81.1	*1929~	USGS	12/22/55	16.17	12,600	1/20	2000	9.52	5,220
Dry Creek near Napa	17.4	1951-	USGS	2/24/58	8.11	3,460	1/20	1800	5.57	1,140
Napa River near Napa	218	*1929-	USGS	1/31/63	27.59	16,900	1/20	2300	16.15	5,260
Redwood Creek										

TABLE A-1 (Continued)

Channe and Chat	Drainage	Period	Source of	Pre	vious Maxi of Record			1963 - 1964			
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Diachg. in cfs	
San Francisco Bay Area	(Continued)										
San Ramon Creek at San Ramon	5.89	1952-	USGS	10/13/62	16.98	1,600	1/20	2100	5-57	480	
San Ramon Creek at Walnut Creek	50.8	1952-	USQS	1/31/63 12/23/55	14.40 14.55	7,980 6,890	1/20	2300	7.82	2,130	
Walnut Creek at Walnut Creek	79.2	1952-	USGS	4/2/58	20.2	12,200	1/20	2200	6.95	2,760	
San Lorenzo Creek at Hayward	37.5	*1939-	USGS	10/13/62 12/22/55	19.73 ^h 20.82 ^h	7,460	1/20	2200	8.58	768	
Arroyo Mocho nr. Pleasanton	143	1962-	USGS	2/1/63	8.60	1,760	1/21	0100	3.07	200	
Arroyo Valle nr. Livermore	147	*1912-	USGS	12/23/55	13.93 ^h	18,200	1/21	0600	3.74	288	
Arroyo Valle at Pleasanton	171	1957-	USGS	4/3/58	25.36	11,300	1/23	1000	7.66	217	
Alameda Creek near Niles	633	1891-	USGS	12/23/55	14.9	29,000 ^c	1/22	1800	5.89	1,850	
Patterson Creek at Union City	-	1958-	USGS	2/1/63	20.4 ^h	10,500	1/22	2100	10.46	1,100	
Alameda Creek at Union City	653	1958-	USGS	2/1/63	19.25 ^h	1,770	1/22	2130	11.11	187	
Coyote Creek near Madrone	196	*1902-	USGS	3/7/11	-	25,000		N	o Peak ^c		
Upper Penitencia Creek at San Jose	21.5	1961-	USGS	3/28/63	3.53	295	1/21	1930	3.98	88	
Alamitos Creek nr. New Almaden	31.9	1958-	USGS	4/2/58	9.67	4,300	1/20	2200	5.00	710	
Los Gatos Cr. at Los Gatos	38.6	*1929-	USGS	2/27/40	14.71 ^b	7,110	1/20	1900	5.00	127	
Guadalupe River at San Jose	146	1929-	USGS	4/2/58	16.55	9,150 ^c	1/20	2400	5.10	1,980	
Saratoga Creek at Saratoga	9.22	1933-	USGS	12/22/55	6.40	2,730	1/20	2130	3.85	338	
Matadero Creek at Palo Alto	7.24	1952-	USGS	12/22/55 12/23/55	9.60 9.88k	854 -	1/20	2030	2.34	223	
San Francisquito Creek at Stanford University	37.5	*1930 -	USGS	12/22/55	13.60	5,560	1/20	2200	4.50	785	
Redwood Creek at Redwood City	1.82	1959 -	USGS	1/31/63	9.36	644	1/20	2200	4.14	114	
Pescadero Creek near Pescadero	45.9	1951-	USGS	12/23/55	21.27	9,420	1/20	2300	9.06	1,170	

TABLE A-1 (Continued)

	Drainage	Period	Source	Pre	vious Maxi			1963 - 1964				
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs		
Central Coastal Area												
San Lorenzo River at Big Trees	111	1936-	USGS	12/23/55	22.55	30,400	1/21	0100	7.29	2,660		
Branciforte Creek at Santa Cruz	17.3	1940-43 1952-	USGS	12/22/55	22.04	8,100	1/20	2300	9.05	903		
Soquel Creek at Soquel	40.2	1951-	USGS	12/23/55	22.33	15,800	1/20	2400	7.55	1,390		
Llagas Creek nr. Morgan Hill	19.6	1951-	USGS	4/2/58	8.45	3,190 ^c	1/20	2200	1.37	21°		
Bodfish Creek near Gilroy	7.40	1959-	USGS	1/31/63	8.25	1,240	1/22	0100	4.87	236		
Tres Pinos Creek near Tres Pinos	206	1939-	USGS	4/4/41	7.75	8,060	1/23	1100	4.53	43		
San Benito River near Hollister	586	1949-	USGS	4/3/58	16.30	11,600	1/24	0100- 0400	3.11	44°		
Pajaro River at Chittenden	1,186	1939-	USGS	12/24/55 4/3/58	32.46 3 3.11	24,000	1/22	0900	9.24	1,460		
Corralitos Creek near Corralitos	10.6	1957-	USGS	4/2/58 1/31/63	7.55 7.62	1,970 1,920	1/20	2200	3.74	299		
Corralitos Creek at Freedom	27.8	1956-	USGS	12/22/55	15.6 ^h	3,620	1/20	2400	5.54	702		
Salinas River near Pozo	74.1	1942-	USGS	1/21/43	13.35	7,210	1/21	0130	3.32	8.2		
Salinas River above Pilitas Creek nr. Santa Margarita	114	1942-	USGS	4/3/58	8.68	4,720 ^c	1/22	0200	1.12	6.5°		
Jack Creek nr. Templeton	25.3	1949-	USGS	1/25/56	9.56	5,040	1/21	0100	4.34	352		
Salinas River at Paso Robles	389	1939-	USGS	3/9/43	16.2	14,200 ^c	1/22	1130	8.12	237°		
Estrella River near Estrella	922	1954-	USGS	4/6/58	7.20	8,850	1/23	1700	2.09	12		
Nacimiento River near Bryson	140	1955-	USGS	12/23/55	24.63	30,300	1/21	0030	11.71	6,070		
San Antonio River at Pleyto	284	*1922-	USGS	4/3/58	6.44	19,100	1/22	1330	2.70	624		
Salinas River near Bradley	2,535	1948-	USGS	4/3/58	12.53	28,400 ^c	1/23	0600	4.48	595 ^c		
Arroyo Seco near Soledad	244	1901-	USGS	4/3/58	14.40 ^b	28,300	1/21	0330	10.34	4,850		

TABLE A-1 (Continued)

	Drainage	Period	Source	Pre	vious Maxi		1963 - 1964				
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs	
Central Coastal Area	(Continued)										
Salinas River near Spreckels	4,156	*1900-	USGS	2/12/38 1/16/52	25.0 26.85	75,000 ^d	1/23	0900	10.25	609 ^c	
Big Sur River near Big Sur	46.5	1950-	USGS	4/2/58	11.56	5,680	1/20	2300	6.48	1,470	
Arroyo De La Cruz near San Simeon	41.4	1950-	USGS	12/23/55	12.40	17,700	1/21	0030	6.72	2,700	
Santa Rosa Creek near Cambria	12.5	1957-	USGS	2/1/60 12/?/55	10.36 15.2 h	2,520	1/20	2400	4.27	246	
Arroyo Grande at Arroyo Grande	102	1939-	USGS	1/15/52	11.97	5,370	1/21	0100	1,46	9.1	
Sisquoc River near Garey	472	1940-	USGS	1/23/43	8.46 ^b	13,000		No Flo	ow During	Year ^c	
Santa Maria River at Guadalupe	1,742	1940-	USGS	1/16/52	8.18	32,800		No Fl	ow During	Year ^C	
Santa Ynez River below Gibraltar Dam, nr. Santa Barbara	216	1920-	USGS	3/2/38	-	35,500°	1/22	0130	4.61	3.2°	
Santa Cruz Creek near Santa Ynez	73.9	1941-	USGS	4/3/58	10.27	3,580	1/22	1300	3.50	22	
San Jose Creek near Goleta	5.51	1941-	USGS	1/21/43	12.74	1,960	1/21	0330	2.96	84	
Atascadero Creek near Goleta	18.3	1941-	USGS	1/15/52	10.85	4,500	1/21	0200	6.98	148	
Carpinteria Creek near Carpinteria	13.1	1941-	USGS	1/15/52	9.75	2,440	1/22	0900	3.50	10	
Central Valley Area											
Sacramento River at Delta	427	1944-	USGS USBR	12/22/55	19.50	37,000	1/20	1300	12.00	13,100	
N. F. Pit River near Alturas	209	1929 - 32 1957 -	USGS	10/14/62	11.07	2,530	1/21	0700	2.15	142	
Pit River nr. Bieber	2,970	*1904-	USGS	3/19/07	16.7	33,800	1/21	1600	4.84	915	
Pit River below Pit No. 4 Dam	4,860	1922-	USGS	12/12/37	17.90	30,200	1/20	1800	4.62	128	
Pit River near Montgomery Creek	5,170	1944-	USGS	12/23/55	14.12	37,100		B	ackwater		
Squaw Creek above Shasta Lake	65.3	1944-	USGS USBR	12/21/55	21.90	17,800	1/20	1400	17.50	8,750	

TABLE A-1 (Continued)

	Drainage	Period	Source of	Pre	vious Maxi of Record		1963 - 1964			
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
			.,,							
Central Valley Area (C	ontinued)									
McCloud River above				((
Shasta Lake	606	1945-	USGS USBR	12/22/55	28.20	45,200	1/20	1500	19.53	13,090
Secure and a Division										
Sacramento River at Keswick	6,710	1938-	USGS DWR	2/23/40	47.2 ^b	186,000	1/20	1230	11.80	8,020
			Dan							
Clear Creek at French Gulch	115	1950-	USGS	12/22/55	13.49	7,050	1/20	1300	9.81	3,300
				, , ,						
Clear Creek near Igo	228	1940-	USGS	12/21/55	13.75	24,500	1/20	1230	6.28	3,030
Cow Creek near Millville	425	1949-	USGS	12/27/51	21.55	45,200	1/20	2000	14.25	17,900
Cottonwood Creek										
near Cottonwood	922	1940-	USGS	3/1/41	15.4	52,300	1/20	2000	13.25	13,000
Battle Creek below										
Coleman Fish Hatchery near Cottonwood	358	1961-	USGS	12/11/37	15.8 ^{h,b}	35,000	1/20	1830	10.99	7,270
Paynes Creek nr. Red Bluff	92.7	1949-	USGS	12/1/61	11.33	10,600	1/20	2030	8.40	4,430
Garage Blanco										
Sacramento River near Red Bluff	9,300	1892-	USGS	2/28/40	38.9	291,000	1/21	0200	14.80	61,600 ^c
Sacramento River										
at Red Bluff	-	-	DWR	2/28/40	32.2	-	1/21	0200	18.27	61,500 ^c
Red Bank Creek				- 1 15-	0.6-		. /25	7 C lan	6.00	7 1000
near Red Bluff	93.5	1959-	DWR USBR	1/31/63	8.67	5,770	1/20	1640	6.82	1,420
A. 4 3										
Antelope Creek near Red Bluff	123	1940-	USGS USCE	2/22/56	12.43	11,500	1/20	1830	12.09	7,280
			0501							
Elder Creek near Paskenta	95.8	1948-	USGS	2/24/58	13.90	11,700	1/20	1600	8.11	2,280
				, ,-						
Elder Creek at Gerber	136	1949-	USGS	2/19/58	14.40 ^b	11,000	1/20	1930	8.53	2,600
			USBR							
Mill Creek near	101	*1909-	USGS	12/11/37	23.4 ^h	23,000	1/20	1900	9.82	7,000
Los Molinos	131	-1303-	abau	12/11/31	23.4	23,000	1/20	1900	9.02	1,000
Thomes Creek at Paskenta	194	1920-	USGS	12/21/55	13.89	23,500	1/20	1530	7.79	3,390
	-,	-)	DWR	,, -,			,			
Deer Creek					h	0	. /	. 0		g 00a
near Vina	208	*1911-	USGS DWR	12/10/37	19.2 ^h	23,800	1/20	1800	10.30	7,880
Sacramento River at Vina Bridge	-	1945-	DWR	2/25/58	89.42	147,000	1/21	0730	80.22	68,100
			USBR							

TABLE A-1 (Continued)

	Drainage	Per1od	Source	Pro	evious Max			19	963 - 1964	
Stream and Station	Area in Sq. Mi.	of Record	of Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
Central Valley Area	(Continued)									
Sacramento River at Hamilton City	-	1945-	DWR USBR	12/11/37	150.7	350,000	1/21	1300	39.28	61,600 ^c
Big Chico Creek near Chico	72.5	1930-	usgs	12/10/37	16.6 ^b	8,260	1/20	2000	10.12	3,780
Stony Creek near Fruto	599	1901-1912 1960-	USGS	2/2/09	16.3 ^b	36,000	1/20	1900	6.47	3,400
Stony Creek near Hamilton City	777	*1940-	USGS	2/25/58	18.31	39,900°		No	Flow ^C	
Sacramento River at Ord Ferry	-	*1921~	DWR	2/28/40	121.7	370,000	1/21	1600- 1800	109.0	57,500°
Sacramento River at Butte City	-	*1921-	USGS DWR	2/7/42	96.87	170,000°	1/21	2300	84.80	52,400
Moulton Weir Spill to Butte Basln	-	*1935~	DWR	2/20/58 2/26/58	83.66 83.66	36,000 ^d 36,000 ^d		No	Flow	
Colusa Weir Spill to Butte Basin	-	*1935-	DWR	2/8/42	70.40	86,000 ^d	1/22	0840	63.81	9,810
Sacramento River at Colusa	-	1940-	USGS DWR	2/8/42	69.20	49,000°	1/22	1000	61.88	33,300 ^c
Colusa Basin Drain at Highway 20	-	1924-	DWR	2/21/58	51.93	25,400 ^e	1/22	1200	43.35	339
Butte Creek near Chico	147	1930-	USGS	12/22/55	13.35	18,700 ^c	1/20	2000	7.20	4,940 ^c
Butte Slough to Sutter Bypass at Mawson Bridge	-	*1934-	DWR	3/1/40	68.9	210,000	1/24	1430	50.72	-
Sutter Bypass at Long Bridge	-	1914-	DWR	3/1/40	57•7	210,000	1/24	2000	43.86	-
Tisdale Weir Spill to Sutter Bypass	-	1940-	DWR	3/1/40	53.35	25,700 ^d	1/22	1830	47.59	6,390
Sacramento River at Knights Landing	-	1940-	USGS DWR	12/8/42 12/3/60	41.83 ^k 30.31	30,000°	1/23	1300	35.24	22,200
Big Grizzly Creek near Portola	45.5	*1925-	USGS	2/1/63	8.03	4,080	1/20	2000	2.13	3 ^e Affected by ice
Middle Fork Feather River near Clio	686	1925-	USGS	2/1/63	16.19	14,500	1/20	2000	5.30	384
Middle Fork Feather River near Merrimac	1,068	1951-	USGS	2/1/63	21.65	65,400	1/20	2200	7.14	3,510

TABLE A-1 (Continued)

	Drainage	Period	Source of	Pr	evious Max of Recor			1963 - 1964			
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft,	Dischg. in cfs	
Central Valley Area	(Continued)										
South Fork Feather River at Enterprise	132	1911-	USGS	12/22/55	21.60	19,200	1/20	2100	7.55	1,420	
Peather River											
at Bidwell Bar	1,347	*1911-	USGS	12/23/55	25.5	104,000	1/20	2145	10.15	8,530	
North Fork Feather											
Prattville	493	*1905-	USGS	3/19/07	16.2 ^b	10,000	1/20	-	1.86	12°	
Indian Creek near Crescent Mills	739	*1906-	USGS	12/24/55	17.80	31,500°	1/20	2300	5.24	1,050°	
	129	1900-	0000	12/24/33	17.00	51,500	1/20	2500	J. L.	1,000	
Spanish Creek above Blackhawk Creek, at Keddie	184	1933-	USGS	2/1/63	13.37	15,000 ^c	1/20	_	5,83	2,190 ^{h,c}	
·	104	1933-	0000	2/1/03	23.31	1),000	1/20		7,05	2,150	
North Fork Feather River at Pulga (formerly Big Bar)	1,953	*1910~	USGS	12/23/55	35,60	72,400 ^c ,g	1/20	1900	12,69	5,970	
	1,900	"1910-	0505	12/23/33	37,00	12,400	1/20	1900	12.09	5,510	
West Branch Feather River	112	1957-	USGS	1/31/63	23.35	21,200°	1/20	1830	11.65	4,850	
ear Paradise	113	1957 -	0505	1/31/03	23.33	21,200	1/20	1030	11,05	4,050	
eather River t Oroville	3,632	1901~	USGS	3/19/07	39•3 ^b	230,000	1/20	2300	144.40	31,200 ^c	
			DWR								
eather River near Gridley	-	*1929-	DWR	12/23/55	102.25	-	1/21	0340	85.18	23,500	
South Honcut											
reek near Bangor	30.5	1950-	USGS	10/13/62	12.40	8,280	1/20	1945	8.62	2,960	
Feather River											
at Yuba City	-	1944-	DWR	12/24/55	82.42	-	1/22	0310	53.23	21,500	
Middle Yuba River above											
Oregon Creek	162	1940-	USGS	1/31/63	18.55	31,600°	1/20	2200	6.09	1,680°.	
Oregon Creek near North											
San Juan	34.4	1911-	USGS	12/22/55	11.90	5,390	1/20	2100	5.97	800	
North Yuba River below											
Goodyears Bar	250	*1930-	USGS	2/1/63	23.8	40,000	1/20	2200	6.06	1,720	
Worth Yuba											
River below Bullards Bar Dam	487	1940-	USGS	1/31/63	42.0 ^h	83,000°	1/21	0700	8.90	2,060 ^c	
				,							
South Yuba River nr. Cisco	51.8	1942-	USGS	1/31/63	20.6 ^h	18,400 ^c	1/22	1230	2,55	48 ^c	
South Yuba											
River at Jones Bar	310	1940-48 1959-	USGS	1/31/63	21.5 ^h	40,000	1/21	0100	9.02	2,500	
		JJJ									

TABLE A-1 (Continued)

	Drainage	Period	Source	Pre	of Record			19	63 - 1964	
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Diachg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
Central Valley Area	(Continued)									
Yuba River at Englebright Dam	1,104	1941-	USGS PGE	2/1/63	544.84	150,000 ^{c,f}	1/20	2300	529.98	7,760 ^c
Deer Creek near Smartville	84.6	1935-	USGS	10/13/62	13.77	11,600°	1/20	2100	10.36	6,170 ^c
Yuba River near Marys v ille	1,340	*1940-	USGS	12/23/55	88.85	160,000°	1/21	0100	68.07	16,300°
Bear River near Auburn	139	1940-	USGS	12/22/55	16.56 ^b	19,700	1/21	0200	9.58	3,900
Bear River nr. Wheatland	292	1928-	USGS	12/22/55	19.30	33,000°	1/21	2100	2.86	1,260 ^c
Feather River at Nicolaus	5,928	1943-	USGS DWR	12/23/55	51.60	357,000 ^c	1/22	1200	37 .7 8	34,000°
Sacramento River at Fremont Weir (West End)	-	*1935-	DWR	12/23/55	39.72	293,800 ^d	1/23	1610	33.47	No Flow Over Weir
Sacramento River at Verona	-	1929-	USGS DWR	3/1/40	41.20	79,200 ^c	1/23	1000	31.36	50,500
Sacramento Weir Spill to Yolo Bypass, near Sacramento		*1939-	USGS DWR	3/26/28 12/23/55	31.83 33.01	118,000 ^d			No Flow	
North Fork American River at North Fork Dam	343	1941-	USGS	1/31/63	11.30	59,700 ^c	1/21	0230	3.29	3,740 ^c
Rubicon River near Foresthill	311	1958-	USGS	2/1/63	35.0 ^h	83,000°	1/21	0600	10.60	990°
Middle Fork American River near Auburn	612	1911-	USGS	2/1/63	43.1 ^h	121,000	1/21	0500	9.27	3,410
South Fork American River near Kyburz	193	1907, 1922-	USGS	2/1/63	10.53	15,600 ^{c,g}	1/20	2330	2.12	66°
South Fork American River near Camino	493	1922-	USGS PGE	12/23/55	32.6 ^h	49,800 ^c	1/21	0330	6.42	1,590 ^c
South Fork American River near Lotus	673	1951-	USGS	12/23/55	21.37	71,800°	1/20	2230	8.31	4,000 ^c
American River at Fair Oaks	1,889	1904-	USGS	11/21/50	31.85 ^b	180,000	1/21	2300	2.93	3,000

TABLE A-1 (Continued)

Stream and Station	Drainage Area in	Period	Source	Pre	vious Max of Recor			19	963 - 196	4
Stream and Station	Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
Central Valley Area	(Continued)									
Sacramento River at Sacramento	-	*1879-	USGS DWR	11/21/50	30.14 ^b	104,000 ^C	1/23	1630	17.31	52,800 ^e
Sacramento River at Walnut Grove	-	1929-	DWR	11/21/50	13.0	-	1/23	1210	7.18	-
Adobe Creek nr. Kelseyville	6.39	1954-	USGS	1/31/63	9.22	1,450	1/20	1400	7.50	800
Kelsey Creek nr. Kelseyville	37.2	1946-	USGS	12/21/55	12.80	8,800	1/20	1600	10.95	5,310
Cache Creek near Lower Lake	528	1944-	USGS	2/24/58	9.40	8,000°		Regulat	ted - Low	Flow
North Fork Cache Creek near Lower Lake	198	1930-	USGS	12/11/37	13.98 ^h	20,300	1/20	1900	8.55	6,020
Cache Creek near Capay	1,052	1942-	USGS	2/24/58	20.90	51,600°	1/21	0430	10.13	7,710 ^c
Cache Creek at Yolo	1,137	1903-	USGS	2/25/58	33.11	41,400 ^{c,g}	1/21	1000	13.17	7,340 ^{c,g}
Yolo Bypass near Woodland	-	1939-	USGS DWR	2/8/42	32.00	272,000			No Flow	
Dry Creek near Middletown	8.41	1959-	USGS	2/8/60	9.90	3,470	1/20	1800	7.52	1,160
Putah Creek near Winters	577	1930-	USGS DWR	2/27/40	30.5	81,000	1/20	5000	6.18	252 ^c
Yolo Bypass near Lisbon	-	1914-	DWR	12/24/55	23.4 ^b	304,800	1/24	1700	9.69	Flow Confined to Low Water Channel
Sacramento River near Rio Vista	-	1906-	USCE DWR	12/25/55	10.2				Tidal	
North Fork Cosumnes River near El Dorado	205	1911 - 41 1948-	USGS	12/23/55	14.8	15,800 ^c	1/22	0430	4.77	598 ^c
Middle Fork Cosumnes River near Somerset	107	1957-	USGS	2/1/63	16.20	11,800	1/21	0130	5.80	309
South Fork Cosumnes River near River Pines	64.3	1957~	USGS	2/1/63	10.90	5,540	1/21	0130	3.38	565

TABLE A-1 (Continued)

	Drainage	Period	Source of	Previous Maximum of Record			1963 - 1964			
Stream and Station	Area in Sq. Mi.	of Record	Record (a)	Date	Stage in ft.	Dischg. in cfs	Date	Time	Stage in ft.	Dischg. in cfs
Central Valley Area	(Continued)									
Cosumnes River at Michigan Bar	537	1907-	USGS DWR	12/23/55	14.59	42,000°	1/22	0700	6.29	4,010 ^c
Cosumnes River at McConnell	724	1941-	USGS USBR DWR	12/23/55	46.26	54,000°	1/22	1400	39.28	5,470 ^c
Dry Creek near Galt	329	1926 - 1933 1944-	USGS USBR DWR	4/3/58	15.28	24,000	1/21	0100	11.30	1,630

LEGEND

(a) USCE - United States Corps of Engineers
USGS - United States Geological Survey
USBR - United States Bureau of Reclamation
DWR - Department of Water Resources
PGE - Pacific Gas and Electric Company
b - Site and datum then in use
c - Affected by storage and/or diversion
d - Discharge over weir
e - Estimated
f - Includes flow through powerhouse
g - Includes flow bypassing station
h - From flood marks
k - Discharge not determined; affected by backwater
l - Crest stage gage
* Incomplete Record


** Maximum of Record













END

AGING STATION
ECIPITATION STATION

AKE

SASIN BOUNDARY

OF RAINFALL IN INCHES

PERIOD JAN. 15-25, 1964



+380

STATE OF CALIFORNIA
THE RESOURCES AGENCY

DEPARTMENT OF WATER RESOURCES
DIVISION OF OPERATIONS

HIGH WATER OF 1963 - 64
JANUARY 1964 STORM

NORTH COASTAL AREA
AL AND STATION LOCATION MAP

NORTH COASTAL AREA STREAM GAGING STATIONS

- 1 Middle Fork Smith River at Gasquet 2 Smith River near Crescent City
- 4 Scott River near Fort Jones
- 5 Klamath River near Seind Valley

14 North Fork Trinity River at Helena

17 South Fork Transty River at Forest Gleno

18 South Fork Trinity River near Hyampom

15 Trinity River near Burnt Ranch 16 New River at Denny

- 7. North Fork Salmon River near Forks of Salmon 8 Salmon River at Somesbar
- 9 Klamath River at Somesbar
- 10. Red Cap Creek sear Orleans
- 1] Bluff Creek near Westchpec
 - 56 Russian River near Hopland
 - 57 Feliz Creek near Hopland

 - 62 Dry Creek near Geyserville
 - 63 Santa Rosa Creek near Santa Rosa
- 14 Hayfork Creek near Restork
- 20 Hayfotk Creek near Hyampom
- 23 Trinsty River near Hoops
- 25 Redwood Creek at Orick
- 26 Little River of Crannell
- 27 Nad River near Forest Glena
- 28 North Fork Mad River near Korbel
- 29 Mad River near Arcute 30 Taroby Creek oper Freehwate

- 32 Eel River below Scott Dam newr Potter Valley
- 33 Eel River at Van Arsdale Dam, near Potter Valley
- 34 Outlet Creek near Longvale
- 35 Eel River above Dos Rius
- 36 Black Butte River near Covel-
- 17 Middle Fark Eel River below Black Butte. Raver, near Covelo
- 38 Eel River below Dos Rips
- 39 North Fork Eel River near Mina
- 40 Eel River at Alderpoint
- 41. South Fork Eel River near Brunscomb
- 42 Tenmile Creek near Laytonville
- 43. South Fork Eel River near Mirando
- 44 Bull Creek near Woott
- 45 Lorabee Creek ness Holmes
- 46 Eel River at Scotta

NORTH COASTAL AREA STREAM GAGING STATIONS (CONTINUED)

- 47 South Fork Van Duzen River near Bridgeville
- 48 Van Duzen River near Bridgeville
- 49 Mattole River near Petrolin
- 50 Noyo River near Fort Bragg
- 51 Rancheria Creek near Boonville 52 Navarro River near Navarro
- 53 South Fork Guelala River near Annapolis
- 54 Russian River near Ukinh
- 55 East Fork Russian River near Calcella


- 58 Russian River near Cloverdale
- 59 Big Sulphur Creek agar Cloverdale
- 60. Russian River near Healdaburg
- 61. Dry Creek near Cloverdale
- 64 Russian River near Guerneville
- b5 Austin Creek near Cazadero

NORTH COASTAL AREA HOURLY PRECIPITATION STATIONS

- Crescent City Maintenance Station
- 2 Happy Camp Ranger Station
- 3 Klamath

- 7 Eureka WB City
- 8 Kneeland 10 SSE
- 10 Miranda Spengler Runch
- 11 Lake Mountain
- 12 Covelo Eel River Ranger Station

- 14 Fort Bragg
- 15 Willits Howard Forest Runger Station
- 16 Redwood Valley
- 17 Navarro I NW
- 18 Point Arena
- 19 The Geysers 20 Venado



SAN FRANCISCO BAY AREA STREAM GAGING STATIONS

- 1 Walker Creek near Tomales
- 2 Corte Medera Creek at Ross
- 3 Noveto Creek near Noveto
- 4 Sonoma Creek at Boyes Hot Springs
- 5 Napa River neer St. Helena
- b Dry Creek near Napa
- 7. Napa River near Napa
- 8 Redwood Creek near Napa
- 9 San Ramoo Creek at Sen Remon
- 10. San Ramon Creek at Walnut Creek
- 11. Walnut Creek at Walnut Creek
- 12 San Lorenzo Creek at Hayward
- 13. Arroyo Mocho near Pleesanton
- 14 Arroyo Valle cear Livermore 15 Arroyo Valle at Pleasanton
- 16 Alamede Creek near Niles
- 17 Patterson Creek at Union City
- 18 Alameda Creek at Union City
- 19 Coyote Creek near Madrone
- 20. Upper Penitencia Creek at San Jose
- 21 Alamitos Creek near New Almaden
- 22. Los Gatos Creek at Los Gatos
- 23 Gundalupe River at San Jone
- 24 Saratoga Creek at Saratoga
- 25 Matadero Creek at Palo Alto
- 26. San Francisquito Creek at Stanford
- 27 Redwood Creek at Redwood City
- 28 Pesendero Creek near Pescadero

SAN FRANCISCO BAY AREA HOURLY PRECIPITATION STATIONS

- 1 St Helena 4 W5W
- 2 Petaluma 1 N
- 3 Noveto 8 WNW
- 4 Mount Temelpais 2 SW
- 5 Martines 3 S
- 6 Walnut Creek 2 ENE
- 7 Oakland WB AP
- 8 San Francisco FOB
- 9 San Francisco WB AP
- 10 SE Faralion
- 11 Hayward 6 ESE
- 12 Palo Alto City Hall
- 14 Morgan Hill 6 WNW



CENTRAL COASTAL AREA STREAM GAGING STATIONS


- 1 San Lozenzo River at Big Trees
- 2 Branciforte Creek at Santa Cruz
- 3. Saquel Creek at Soquel
- 4 Lingus Creek near Morgan Hill
- 5 Bodfish Creek near Gilroy
- 5. Tres Pinos Creek near Tres Pinos 7. San Benito River near Hollister
- 8. Pajaro River at Chittenden
- 9. Corralitos Creek near Corralitos
- 10 Constitut Creek at Freedom
- 11 Salinus River near Pozo
- 12 Salinas River above Pilitas Creek near Santa Marganta
- 13 Jack Creek near Templeton
- 14. Salinas River at Paso Robles
- 15. Estrella River near Estrella
- 16 Nacimiento River near Bryson 17 San Antonio River at Pleyto
- 18. Salines River near Bradley
- to settings have bear bin
- 19. Arroya Seca near Saledad
- 20 Satinas River near Spreckets 21 Big Sur River near Big Sur
- 22. Amoyo De Lu Cruz near San Simeon
- 23 Santa Rosa Creek near Cambria
- 24 Arroyo Grande el Arroyo Grande
- 25 Sisquot River pear Garey
 26 Santa Maria River at Guadalupe
- Santa Ynez River below Gibraltar Dam. near Santa Barbara
- 28 Santa Cruz Creek near Santa Ynez
- 29 San Jose Creek near Goleta

CENTRAL COASTAL AREA STREAM GAGING STATIONS ICONTINUEDI

- 30 Atascadero Creek near Goleta
- 31 Carpinteria Creek near Carpinteria

CENTRAL COASTAL AREA HOURLY PRECIPITATION STATIONS

- I Boulder Creek Locatelli Ranch
- 2 Mount Madonna
- 3 Hollister 10 ENE
- 4. Sun Juan Bautista
- 5 Upper Tres Pinos 6 Del Monte
- 7 Gonzales 9 ENE
- 8 Arroyo Seco
- 9 King City
- 10 Luckwood 2 N
- 11 Bryson
- 12 Paso Robles 5 NW
- 13 La Panza Ranch
- 14. San Luis Obispo Cal Poly
- 15 Cuyuma Ranch 16 Santa Marie WB AP
- 17 Cuyama Ranger Station
- 18 Point Arguello 19 Cachuma Dam
- 20 Santa Berbara



PASTONI ANTINA AREA

STREAM CAGING STATIONS

W. Loctomesty River at Frem at Ba

40 Carbo Lord near Lover Labo

III to ober Konner to A sin Mill Son Appell Hotel Committee Street want his Phone St.


LASTRAL VALLEY OREA


HOLDS A RESCUBLING TON STATIONS

68 Louiser




LEGEND STREAM GAGING STATION MODELL DOCCIOITATION PRESIDE NATURAL LAKE - BRAINAUF BASIN BOUNDARY FOR THE PERIOD JAN 15+25, 1964 DEPARTMENT OF WATER RESOURCES HIGH WATER OF 1963 - 64 JANUARY 1964 STORM CENTRAL VALLEY AREA SAN FRANCISCO BAY AREA ISOHYETAL AND STATION LOCATION MAP











THIS BOOK IS DUE ON THE LAST DATE STAMPED BELOW

RENEWED BOOKS ARE SUBJECT TO IMMEDIATE RECALL

APR 5 1976 APR 5 1976

LIBRARY, UNIVERSITY OF CALIFORNIA, DAVIS

Book Slip-25m-6,'66 (G3855s4)458

Nº 479869

California. Dept. of Water Resources. Bulletin.

PHYSICAL SCIENCES LIBRARY TC821, C2 A2 no.69:61, c.2

LIBRARY UNIVERSITY OF CALIFORNIA DAVIS

Call Number:

California. De.t. of Nater Resources. Bulletin.

TC824 C2 A2 no.69:64



